

Total No. of Questions—24

Total No. of Printed Pages—4

Regd. No.

--	--	--	--	--	--	--	--	--	--

Part III
MATHEMATICS
Paper I-A
(English Version)

Time : 3 Hours**Max. Marks : 75****Note :—** This question paper consists of THREE Sections A, B and C.**SECTION A**

10×2=20

(I) Very short answer type questions :

(i) Answer ALL questions.

(ii) Each question carries TWO marks.

1. If $A = \{-2, -1, 0, 1, 2\}$ and $f : A \rightarrow B$ is a surjection defined by $f(x) = x^2 + x + 1$, then find B.

2. Find the domain of the real valued function $f(x) = \sqrt{4x - x^2}$.

3. If $A = \begin{bmatrix} 2 & -4 \\ -5 & 3 \end{bmatrix}$, then find $A + A'$ and AA' .

4. Find the rank of the matrix $\begin{bmatrix} 1 & 4 & -1 \\ 2 & 3 & 0 \\ 0 & 1 & 2 \end{bmatrix}$.

5. If the position vectors of the points A, B and C are $-2\vec{i} + \vec{j} - \vec{k}$, $-4\vec{i} + 2\vec{j} + 2\vec{k}$ and $6\vec{i} - 3\vec{j} - 13\vec{k}$ respectively and $\overline{AB} = \lambda \overline{AC}$, then find the value of λ .
6. Find the vector equation of the plane passing through the points $\vec{i} - 2\vec{j} + 5\vec{k}$, $-5\vec{j} - \vec{k}$ and $-3\vec{i} + 5\vec{j}$.
7. $\vec{a} = 2\vec{i} - \vec{j} + \vec{k}$, $\vec{b} = \vec{i} - 3\vec{j} - 5\vec{k}$. Find the vector $\vec{c}$ such that $\vec{a}$, $\vec{b}$ and $\vec{c}$ form the sides of a triangle.
8. Sketch the graph of the function $\sin 2x$ in the interval $(0, \pi)$.
9. Evaluate $\cos^2 52\frac{1^\circ}{2} - \sin^2 22\frac{1^\circ}{2}$.
10. If $\sinh x = 3$, then show that $x = \log_e (3 + \sqrt{10})$.

SECTION B

5×4=20

(II) Short answer type questions :

- (i) Answer ANY FIVE questions.
- (ii) Each question carries FOUR marks.

11. If $A = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}$ is non-singular matrix, then A is invertible and

prove that $A^{-1} = \frac{\text{Adj } A}{\det A}$.

12. If $\vec{a}$, $\vec{b}$, $\vec{c}$ are non-coplanar vectors, prove that the following four points are co-planar $6\vec{a} + 2\vec{b} - \vec{c}$, $2\vec{a} - \vec{b} + 3\vec{c}$, $-\vec{a} + 2\vec{b} - 4\vec{c}$, $-12\vec{a} - \vec{b} - 3\vec{c}$.

13. If $\bar{a} + \bar{b} + \bar{c} = 0$, $|\bar{a}| = 3$, $|\bar{b}| = 5$ and $|\bar{c}| = 7$, then find the angle between $\bar{a}$ and $\bar{b}$.
14. If a, b, c are non-zero real numbers and α, β are solutions of the equation $a \cos \theta + b \sin \theta = c$, then show that :
- (i) $\sin \alpha + \sin \beta = \frac{2bc}{a^2 + b^2}$
- (ii) $\sin \alpha \cdot \sin \beta = \frac{c^2 - a^2}{a^2 + b^2}$
15. Solve the equation $\cot^2 x - (\sqrt{3} + 1) \cot x + \sqrt{3} = 0$; $0 < x < \frac{\pi}{2}$.
16. Prove that :

$$\tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{13} - \tan^{-1} \frac{2}{9} = 0.$$

17. Prove that :

$$\frac{1 + \cos(A - B) \cos C}{1 + \cos(A - C) \cos B} = \frac{a^2 + b^2}{a^2 + c^2}.$$

SECTION C

5×7=35

- (III) Long answer type questions :

- (i) Answer ANY FIVE questions.
- (ii) Each question carries SEVEN marks.
18. Let $f : A \rightarrow B$, $g : B \rightarrow C$ be bijections, then prove that $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$.
19. Using mathematical induction, prove that :

$$\left(1 + \frac{3}{1}\right) \left(1 + \frac{5}{4}\right) \left(1 + \frac{7}{9}\right) \dots \dots \dots \left(1 + \frac{2n+1}{n^2}\right) = (n+1)^2.$$

20. Find the value of x if
$$\begin{vmatrix} x-2 & 2x-3 & 3x-4 \\ x-4 & 2x-9 & 3x-16 \\ x-8 & 2x-27 & 3x-64 \end{vmatrix} = 0.$$

21. Examine whether the equations are consistent or inconsistent and if consistent find the complete solution :

$$x + y + z = 6$$

$$x - y + z = 2$$

$$2x - y + 3z = 9.$$

22. $\vec{a}, \vec{b}, \vec{c}$ are three vectors, then prove that :

$$(\vec{a} \times \vec{b}) \times \vec{c} = (\vec{a} \cdot \vec{c}) \vec{b} - (\vec{b} \cdot \vec{c}) \vec{a}.$$

23. If A, B, C are angles in a triangle, then prove that :

$$\sin^2 A + \sin^2 B - \sin^2 C = 2 \sin A \sin B \cos C.$$

24. In ΔABC , show that :

$$(r_1 + r_2) \sec^2 \frac{C}{2} = (r_2 + r_3) \sec^2 \frac{A}{2} = (r_3 + r_1) \sec^2 \frac{B}{2}.$$